

## Pharmacology II

Which of the following is the antidote for the toxin Benzodiazepines?

A

- Flumazenil
- Methylene blue
- Deferoxamine
- Alkalinize urine

Which of the following is the antidote for the toxin Lead?

C

- Naloxone
- Nitrite
- CaEDTA
- Dialysis

Which of the following is the primary site of activity for the drug Warfarin?

B

- Kidney
- Liver
- Blood
- Heart

Lansoprazole is not used in which of the following cases?

D

- Gastritis
- Peptic Ulcers
- Zollinger-Ellison syndrome
- Thalamus hypertrophy

Which of the following drugs is associated with the reaction of Cinchonism?

B

- Valproic acid
- Quinidine
- Isoniazid
- Ethosuximide

Which of the following drugs is associated with the reaction of hepatitis?

C

- Valproic acid
- Quinidine
- Isoniazid
- Ethosuximide

Which of the following drugs is associated with the reaction of Stevens-Johnson syndrome?

D  
Valproic acid  
Quinidine  
Isoniazid  
Ethosuximide

Which of the following drugs is associated with the reaction of Tendon dysfunction?

D  
Digitalis  
Niacin  
Tetracycline  
Fluoroquinolones

A drug ending in the suffix (pril) is considered a \_\_\_\_\_.

B  
H<sub>2</sub> agonist  
ACE inhibitor  
Antifungal  
Beta agonist

A drug ending in the suffix (azole) is considered a \_\_\_\_\_.

C  
H<sub>2</sub> agonist  
ACE inhibitor  
Antifungal  
Beta agonist

A drug ending in the suffix (tidine) is considered a \_\_\_\_\_.

D  
Antidepressant  
Protease inhibitor  
Beta antagonist  
H<sub>2</sub> antagonist

A drug ending in the suffix (navir) is considered a \_\_\_\_\_.

B  
Antidepressant  
Protease inhibitor  
Beta antagonist  
H<sub>2</sub> antagonist

Which of the following drugs is associated with the reaction of extreme photosensitivity?

C  
Digitalis  
Niacin

Tetracycline  
Fluoroquinolones

Which of the following is not related to a drug toxicity of Prednisone?

B  
Cataracts  
Hypotension  
Psychosis  
Acne

Which of the following is not related to a drug toxicity of Atenolol?

B  
CHF  
Tachycardia  
AV block  
Sedative appearance

Which of the following is considered a class IA Sodium Channel blocker?

B  
Mexiletine  
Aminodarone  
Quinidine  
Procainamide

Which of the following is considered a class IA Sodium Channel blocker?

A  
Propafenone  
Disopyramide  
Aminodarone  
Quinidine

Potassium sparing diuretics have the primary effect upon the \_\_\_\_\_ found in the kidney.

D  
Proximal convoluted tubule  
Loop of Henle  
Collecting duct  
Distal convoluted tubule

Which of the following is not directly related to a drug toxicity of Nitroglycerin?

D  
Headaches  
Tachycardia  
Dizziness  
Projectile vomiting

Which of the following is not directly related to a drug toxicity of Ibuprofen?

D

Nausea

Renal dysfunction

Anemia

Muscle wasting